Available online at www.sciencedirect.com
ScienceDirect
Nuclear Physics A 00 (2016) 1–4
Nuclear Physics A
www.elsevier.com/locate/procedia

arXiv:1602.00794v1 [nucl-th] 2 Feb 2016

Causal hydrodynamic fluctuation in Bjorken expansion

K. Nagai[a], R. Kurita[a,b,c], K. Murase[a,b,c], T. Hirano[a]

[a]*Department of Physics, Sophia University, Tokyo 102-8554, Japan*
[b]*Department of Physics, the University of Tokyo 113-0033, Japan*
[c]*Theoretical Research Division, Nishina Center, Riken, Wako 351-0198, Japan*

Abstract

We investigate effects of causal hydrodynamic fluctuations in the longitudinally expanding quark gluon plasma on final entropy distributions in high-energy nuclear collisions.

Keywords: Quark gluon plasma, Relativistic fluctuating hydrodynamics, Bjorken expansion

1. Introduction

In high-energy nuclear collisions, the space-time evolution of the quark-gluon plasma (QGP) is well described by relativistic hydrodynamics. So far, a vast body of analyses have been done to extract the bulk and transport properties of the QGP by using event-by-event viscous fluid-dynamical simulations. In these simulations, initial fluctuation of the transverse profile has been mainly focused on to describe the higher order harmonics of azimuthal momentum distributions of observed hadrons. Moreover, whether hydrodynamic description of the QGP can be applied in small colliding systems such as proton-proton and/or proton (deuteron)-nucleus collisions is one of the open problems for these years.

In this study, instead of the initial fluctuations, we investigate hydrodynamic fluctuations [1, 2, 3, 4] to consistently describe the space-time evolution of the QGP on an event-by-event basis. We focus on the boost-invariant Bjorken expansion [5] within the framework of causal hydrodynamic fluctuations to study the effect of them on final entropy distributions.

Throughout this contribution, we employ natural unit $c = \hbar = k_{\rm B} = 1$.

2. Models

We assume the system created in high-energy nuclear collisions expands in a boost-invariant way in the beam direction. The four velocity in this case can be parametrised as [5]

$$u^\mu = (\cosh\eta_{\rm s}, 0, 0, \sinh\eta_{\rm s}), \tag{1}$$

where $\eta_{\rm s} = \frac{1}{2}\ln[(t+z)/(t-z)]$ is space-time rapidity. We neglect baryon density at relativistic energies. Then hydrodynamic equations reduce to

$$\frac{{\rm d}e}{{\rm d}\tau} = -\frac{e+P}{\tau}\left(1 - \frac{\pi}{sT} + \frac{\Pi}{sT}\right), \tag{2}$$

where τ is proper time, e is energy density, P is hydrostatic pressure, s is entropy density and T is temperature. $\pi = \pi^{00} - \pi^{33}$ is an quantity calculated from shear stress tensor $\pi^{\mu\nu}$ in Bjorken expansion and Π is bulk pressure. Time evolution of these dissipative quantities are described by the constitutive equations. Within causal hydrodynamic fluctuations, these equations in the differential form can be written as

$$\tau_\pi \frac{\mathrm{d}\pi}{\mathrm{d}\tau} + \pi = \frac{4\eta}{3\tau} + \xi_\pi, \tag{3}$$

$$\tau_\Pi \frac{\mathrm{d}\Pi}{\mathrm{d}\tau} + \Pi = -\frac{\zeta}{\tau} + \xi_\Pi. \tag{4}$$

Here η and ζ are viscous coefficients and τ_π and τ_Π are the corresponding relaxation times [6] for shear viscosity and bulk viscosity, respectively. In the relativistic dissipative hydrodynamics, there is an issue on causality within the first order theory. So the relaxation terms in the left hand side of Eqs. (3) and (4) are crucial to obey the causality. ξ_π and ξ_Π are Gaussian white noises for shear stress and bulk pressure, respectively. Statistical properties of these noise terms are determined from transport coefficients through the fluctuation-dissipation relations. In the case of Bjorken expansion, the fluctuation-dissipation relations in the discretised version of the stochastic differential equations, (3) and (4), are

$$\langle \xi_\pi(\tau_i)\xi_\pi(\tau_j)\rangle = \frac{8T\eta\delta_{ij}}{3\Delta\tau\Delta V}, \tag{5}$$

$$\langle \xi_\Pi(\tau_i)\xi_\Pi(\tau_j)\rangle = \frac{2T\zeta\delta_{ij}}{\Delta\tau\Delta V}, \tag{6}$$

$$\langle \xi_\pi\rangle = \langle \xi_\Pi\rangle = 0, \quad \langle \xi_\pi\xi_\Pi\rangle = 0. \tag{7}$$

Here $\Delta\tau$ is time step, $\Delta V = \tau\Delta\eta_s\Delta x\Delta y$ is volume of a fluid element and angle bracket means ensemble average. It is noted that variances of noises are inversely proportional to the volume of a fluid element and, therefore, that the effect of hydrodynamic fluctuations could be important in small colliding systems.

As for the equation of state, we first parametrise entropy density as a function of temperature [7]

$$s(T) = d_\mathrm{H}\frac{4\pi^2}{90}T^3[1 - f(T)] + d_\mathrm{Q}T^3 f(T), \tag{8}$$

$$f(T) = \frac{1}{2}\left[1 + \tanh\left(\frac{T - T_\mathrm{c}}{\Gamma}\right)\right]. \tag{9}$$

The other thermodynamic quantities are calculated from

$$P(T) = \int_0^T s(T')dT', \quad e(T) = Ts(T) - P(T). \tag{10}$$

Parameters $d_\mathrm{H} = 3$ and $d_\mathrm{Q} = 37$ are degrees of freedom in the hadronic matter and the quark gluon plasma, respectively. Transition temperature is $T_\mathrm{c} = 0.17$ GeV and a width of transition is $\Gamma = 0.02T_\mathrm{c}$. The equation of state obtained in this way exhibits the cross-over behaviour.

We employ models for transport coefficients as $\eta/s = 1/4\pi$ [8] and $\zeta/s = 15\left(1/3 - c_\mathrm{s}^2\right)$ [9]. Relaxation times are $\tau_\pi = \tau_\Pi = 3\eta/2p$. It should be emphasised that the purpose of the present study is to demonstrate the importance of hydrodynamic fluctuations for shear and bulk viscosities and the choice of these models is not intended for any specific matter nor situation.

3. Results

In the following, we start from a fixed set of initial conditions. Initial temperature is $T_0 = 0.22$ GeV at initial time $\tau_0 = 1$ fm. At the initial time dissipative quantities are assuemed to be vanishing. As a typical size of one fluid element, we choose $\Delta x = \Delta y = 1$ fm and $\Delta\eta_\mathrm{s} = 1$. Time step in the simulations is $\Delta\tau = 0.1$ fm.

Time evolution of the product of the entropy density, s, and the proper time, τ, is shown in Fig. 1 (a). When one neglects dissipative quantities in Eq. (2), the quantity $s\tau$ is the entropy per unit space-time rapidity and per unit transverse area and a constant of motion for a perfect fluid in one-dimensionally expanding system [5]. On the other hand, $s\tau$ increases monotonically for a viscous fluid (neglecting noise terms in Eqs. (3) and (4)) due to production of entropy. Since thermodynamic force is the inverse of the proper time in the Bjorken expansion, the production of entropy is dominated in the early stage (< 10 fm) and the quantity $s\tau$ saturates in the late stage (∼ 50 fm). We also show two results of causal hydrodynamic fluctuations in Fig. 1 (a). Due to the noise terms in Eqs. (3) and (4), the quantity $s\tau$ is fluctuating during the evolution and final entropy is not the same in these two results even though the initial conditions are common. Interestingly, $s\tau$ can decrease temporarily during the time evolution. This behaviour is explained by the entropy production rate σ in the case of causal hydrodynamic fluctuation. From Eqs. (2) - (4), one derives entropy production rate in a fluid element,

$$\sigma = \frac{\mathrm{d}\Delta S}{\mathrm{d}\tau} = \left(\frac{\pi}{T} - \frac{\Pi}{T}\right)\Delta\eta_{\mathrm{s}}\Delta x\Delta y, \tag{11}$$

$$\Delta S = s\tau\Delta\eta_{\mathrm{s}}\Delta x\Delta y. \tag{12}$$

In the case of causal hydrodynamic fluctuations, shear stress and bulk pressure can take both positive and negative values due to noise terms. As a result, entropy production rate sometimes takes negative values. This is dynamical manifestation of the fluctuation-dissipation relation: Fluctuations make the system deviate from the maximum entropy state by reducing entropy, but dissipations move it back to the state by producing entropy.



Fig. 1. (a) Time evolution of entropy for a perfect fluid (dashed), a viscous fluid (solid) and fluctuating fluids (dotted and dash-dotted). (b) Histogram of the entropy distribution at proper time τ = 50 fm

We analysed two thousand events in causal hydrodynamic fluctuations. Figure 1 (b) shows a histogram of the quantity $s\tau$ at τ = 50 fm. The shape of this entropy distribution looks like a Gaussian. The mean value (28.06(15)/fm^2) is almost the same as the one for the viscous fluid (28.05 /fm^2). When the effect of hydrodynamic fluctuations is sufficiently large, one sees the final entropy being smaller than the initial entropy ($s_0\tau_0$ = 22.49 /fm^2) in a small fraction of the total events in Fig. 1 (b). We emphasise here that this is not in contradiction with the second law of thermodynamics. In fact, one is able to quantify the fraction of net negative entropy production from the fluctuation theorem [10].

To see the volume dependence of hydrodynamic fluctuations, we calculate variance of the final entropy distribution normalised by initial entropy,

$$\frac{\langle(s\tau - \overline{s\tau})^2\rangle}{(s_0\tau_0)^2}. \tag{13}$$

Here $\overline{s\tau}$ is the average of the final entropy. Figure 2 shows a normalised variance of the final entropy distribution as a function of volume (= $\Delta\eta_{\mathrm{s}}\Delta x\Delta y$). As expected from Eqs. (3) and (4), the normalised



Fig. 2. Volume dependence of variance of final entropy normalised by initial entropy. The solid line is the inverse of the volume.

variance is inversely proportional to the volume. Within the present setting of initial conditions, an width of the final entropy distribution is a few percent of the initial entropy in a moderate size of the volume (~ 1 fm^2). The smaller the size of a fluid element is, the more significant the effect of hydrodynamic fluctuations is. One would say that the effect of hydrodynamic fluctuation is important in the small colliding systems such as proton-proton, proton (deuteron)-nucleus and, perhaps, peripheral nucleus-nucleus collisions.

4. Summary

We first formulated relativistic causal hydrodynamics with the effects of thermal fluctuations during the evolution (hydrodynamic fluctuations) in the one-dimensionally expanding QGP. We calculated entropy production within this framework and compare them with that from the conventional viscous fluid-dynamics. Final entropy fluctuates around the mean value due to hydrodynamic fluctuations even when the system starts from the same initial condition in a macroscopic sense. The entropy can temporally decreases during the evolution, which can be understood from the fluctuation theorem in non-equilibrium statistical physics. Hydrodynamic fluctuations should be implemented in the hydrodynamic description of the QGP on an event-by-event basis since, through the fluctuation-dissipation relation, the hydrodynamic fluctuations are always accompanied by the corresponding dissipations in viscous fluid-dynamics. Since multiplicity is almost proportional to the final entropy of the system in high-energy nuclear collisions, the effects of hydrodynamic fluctuations would be important in multiplicity distributions in small colliding system.

This work was supported by JSPS KAKENHI Grant Numbers 12J08554 (K.M.) and 25400269 (T.H.).

References

[1] E. Calzetta, Class. Quant. Grav. **15** (1998) 653.
[2] J. I. Kapusta, B. Muller and M. Stephanov, Phys. Rev. C **85** (2012) 054906.
[3] A. Kumar, J. R. Bhatt and A. P. Mishra, Nucl. Phys. A **925** (2014) 199.
[4] K. Murase and T. Hirano, arXiv:1304.3243 [nucl-th].
[5] J. D. Bjorken, Phys. Rev. D **27** (1983) 140.
[6] W. Israel, Ann. Phys. (N.Y.) **100** (1976) 310.
[7] M. Asakawa and T. Hatsuda, Phys. Rev. D **55** (1997) 4488.
[8] P. Kovtun, D. T. Son and A. O. Starinets, Phys. Rev. Lett. **94** (2005) 111601.
[9] S. Weinberg, Astrophys. J. **168** (1971) 175.
[10] T. Hirano, R. Kurita, K. Murase and K. Nagai, Nucl. Phys. A **931** (2014) 831.

ELSEVIER